

UNIVERSITY OF PUNE
[4362]-220
Electrical/Instrumentation/Computer/I.T.
S. E. Examination - 2013
Engineering Mathematics - III
(2008 Pattern)

Total No. of Questions : 12
[Time : 3 Hours]

[Total No. of Printed Pages :6]
[Max. Marks : 100]

Instructions :

- (1) Answer Q1 or Q2, Q3 OR Q4, Q5 OR Q6, From section I and Q7 OR Q8, Q9 OR Q10, Q11 OR Q12 From section II.
- (2) Answers to the **two sections** should be written in **separate answer-books**.
- (3) Neat diagrams must be drawn wherever necessary.
- (4) Black figures to the right indicate full marks.
- (5) Use of logarithmic tables, slide rule, Mollier charts, electronic pocket calculator and steam tables is allowed.
- (6) Assume suitable data, if necessary.

SECTION-I

Q1. (a) Solve (any three)

[12]

1) $(D^2 - 1)y = \cos x \cosh x$

2) $(D^2 + 2D + D)y = e^{-x} \log x$

3) $\frac{d^2y}{dx^2} + \frac{1}{x} \frac{dy}{dx} = 2 + \log x$

4) $\frac{dx}{xy^3 - 2x^4} = \frac{dy}{2y^4 - x^3y} = \frac{dz}{9z(x^3 - y^3)}$

Q.1 (b) An inductor of 0.25 henries is connected in series with a capacitor of 0.04 farads and a generator having alternative voltage given by $12\sin 10t$. Find the charge and current at any time t. [5]

OR

Q2. (a) Solve: (any three) [12]

(1) $(D^2 + 1)y = x \cos 2x$

(2) $(D^2 - 2D + 2)y = x^2 + e^{-x}$

(3) $(D^2 - 2D)y = e^x \sin x$ (variation of parameters)

(4) $((2x + 5)^2 \frac{d^2y}{dx^2} + 8y - 4(2x + 5) \frac{dy}{dx} = 5 \log(2x + 5)$

Q2. (b) Solve: [5]

$$\frac{dx}{dt} + \frac{dy}{dt} - 3x - y = e^t; \frac{dx}{dt} + 2x + y = 0$$

Q3. (a) If $f(z)$ is analytic, prove that $(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}) |f(z)|^2 = 4 |f'(z)|^2$ [5]

(b) Show that the transformation $w = z + \frac{1}{z} - 2i$ maps the circle $|z|=2$ an ellipse. [5]

(c) Evaluate: $\oint_c \frac{z+4}{z^2+2z+5} dz$ where $c: |z+2i| = \frac{3}{2}$ [6]

OR

Q4. (a) If $f(z) = u+iv$ is analytic function, find $f(z)$ if $u+v=3(x + y) + \frac{x-y}{x^2+y^2}$ [5]

(b) Find the bilinear transformation which maps the points 0,1,2 of z -plane to the points $1, \frac{1}{2}, \frac{1}{3}$ of w plane respectively.

(c) Evaluate:

$$\int_0^{2\pi} \frac{d\theta}{5-3\cos\theta}$$
 [6]

Q5. (a) Find fourier transform of

$$f(x)=\begin{cases} \cos x + \sin x & |x| \leq \pi \\ 0 & |x| > \pi \end{cases}$$
 [5]

(b) using fourier integral representation, show that :

$$\frac{2}{\pi} \int_0^{\infty} \frac{(\lambda^2+2)\cos\lambda x}{\lambda^4+4} d\lambda = e^{-x} \cos x, x > 0 \quad [6]$$

(c) find z-transform of (any two) [6]

1) $f(k) = \frac{\sin ak}{k}, k \geq 0$

2) $f(k) = k^2, k \geq 0$

3) $f(k) = \begin{cases} 7^k & k < 0 \\ 5^k & k \geq 0 \end{cases}$

OR

Q6. (a) Find inverse Z-transform of: (Any two) [8]

- 1) $\frac{2z^2-10z+13}{(z-3)^2(z-2)} \quad 2 < |z| < 3$
- 2) $\frac{z(z+1)}{z^2-2z+1} \quad |z| > 1$
- 3) $\frac{z^2}{z^2+4} \quad \text{inversion integral method}$

b) Solve:

$$f(k) - 4f(k-2) = \left(\frac{1}{2}\right)^k, k \geq 0 \quad [4]$$

c) Solve integral equation:

$$\int_0^{\infty} f(x) \sin \lambda x dx = \frac{e^{-a\lambda}}{\lambda}, \lambda > 0 \quad [5]$$

SECTION II

Q.7 (a) Following are the marks of ten students in math's- III and strength of material (SOM) calculate the coefficient of correlation. [8]

M-III	23	28	42	17	26	35	29	37	16	46
SOM	25	22	38	21	27	39	24	32	18	44

(b) Calculate the first four central moments and β_1, β_2 for the following distribution. [9]

x	0	1	2	3	4	5	6	7	8
F	1	8	28	56	70	56	28	8	1

OR

Q8. (a) The mean and variance of Binomial distribution are 6 and 2 respectively

Find: 1) $p(r \leq 1)$ 2) $p(r \geq 2)$ [6]

(b) If the probability that an individual suffers a bad reaction from a certain injection is 0.001, then determine the probability that out of 2000 individuals

- 1) Exactly 3 will suffer a bad reaction
- 2) More than 2 will suffer a bad reaction [6]

(c) A manufacturer of envelopes knows that the weight of envelope is normally distributed with mean 1.9 gm and variance 0.01gm. find how many envelopes weighing

- 1) 2 grams or more
- 2) 2.1 grams or more

Can be expected in a given packet of 1000 envelopes (Given Area for $z=1$ is 0.3413 and Area for $z=2$ is 0.4772) [5]

Q.9 (a) If $\vec{r}(t) = t^2\vec{i} + t\vec{j} - 2t^3\vec{k}$ then [5]

Evaluate $\int_1^2 \vec{r} \times \frac{d^2\vec{r}}{dt^2} dt$

(b) Prove the following (any two) [6]

$$1) \vec{b} \times \nabla[\vec{a} \cdot \nabla \log r] = \frac{\vec{b} \times \vec{a}}{r^2} - 2 \frac{(\vec{a} \cdot \vec{r})(\vec{b} \cdot \vec{r})}{r^4}$$

$$2) \nabla^2 \left(\frac{\vec{a} \cdot \vec{b}}{r} \right) = 0$$

$$3) \nabla \times \left(\frac{\vec{a} \times \vec{r}}{r} \right) = \frac{\vec{a}}{r} + \frac{(\vec{a} \cdot \vec{r})\vec{r}}{r^3}$$

Q9. (c) Find the directional derivative of $\phi = 4xz^3 - 3x^2y^2z$ at (2,-1,2) in direction towards the point (2,-2,4) [5]

OR

Q10. (a) Verify whether $\vec{F} = (y \sin z - \sin x)\vec{i} + (x \sin z + 2yz)\vec{j} + (xy \cos z + y^2)\vec{k}$ is irrotational and if so find the scalar ϕ such that $\vec{F} = \nabla\phi$ [5]

(b) If \vec{u} and \vec{v} are irrotational vectors then prove that $\vec{u} \times \vec{v}$ is solenoidal vector. [5]

(c) If directional derivative of $\phi = ax^2y + by^2z + cz^2x$ at (1,1,1) has maximum magnitude 15 in the direction parallel to $\frac{x-1}{2} = \frac{y-3}{-2} = \frac{z}{1}$
Then find values of a,b,c. [6]

Q11. (a) Find the work done in moving the particle long the curve $x = a \cos\theta$, $y = a \sin\theta$, $z = b\theta$ from $\theta = \frac{\pi}{4}$ to $\theta = \frac{\pi}{2}$ under the field of force given by $\vec{F} = -3a \sin^2\theta \cos\theta \vec{i} + a(2 \sin\theta - 3 \sin^3\theta)\vec{j} + b \sin 2\theta \vec{k}$ [5]

(b) Evaluate $\int \int_S (\nabla \times \vec{F}) \cdot \hat{n} ds$ where [6]

$\vec{F} = (x^3 - y^3)\vec{i} - xyz\vec{j} + y^3\vec{k}$ And S is the surface $x^2 + 4y^2 + z^2 - 2x = 4$ above the plane $x=0$.

(c) Evaluate $\int \int_S (x^3\vec{i} + y^3\vec{j} + z^3\vec{k}) \cdot d\vec{s}$ where S is the surface of the sphere $x^2 + y^2 + z^2 = 16$ [6]

Q.12 (a) Evaluate $\int \int_s \frac{\bar{r}}{r^3} \cdot \hat{n} ds$ by using Gauss Divergence theorem [5]

(b) Use Stoke's theorem to evaluate [6]

$\int_c (4y\bar{i} + 2z\bar{j} + 6y\bar{k}) \cdot d\bar{r}$ where 'c' is the curve of intersection of $x^2 + y^2 + z^2 = 2z$ and $x = z - 1$

(c) Two of the maxwell's equation are $\nabla \cdot \bar{B} = 0$, $\nabla \times \bar{E} = -\frac{\partial \bar{B}}{\partial t}$. given $\bar{B} = \text{curl } \bar{A}$ then deduce that $\bar{E} + \frac{\partial \bar{A}}{\partial t} = -\text{grad}(v)$ where V is a scalar point function.