

Total No. of Questions : 12]

SEAT No. :

P2020

[Total No. of Pages : 6

F.E. (Semester - II)
ENGINEERING MATHEMATICS - II
(2008 Course)

Time : 3 Hours]

[Max. Marks : 100

Instructions to the candidates:

- 1) *In section - I Solve Q.1 or Q.2, Q.3 or Q.4, Q.5 or Q.6. In section-II Solve Q.7 or Q.8, Q.9 or Q.10, Q.11 or Q.12.*
- 2) *Neat diagrams must be drawn wherever necessary.*
- 3) *Figures to the right indicate full marks.*
- 4) *Use of non - programmable electronic pocket calculator is allowed.*
- 5) *Assume suitable data, if necessary.*

SECTION - I

Q1) a) Form the differential equation whose general solution is
 $y = ae^{2x} + be^{3x}$. [6]

b) Solve any two [10]

i) $\frac{dy}{dx} = \frac{y+2}{x+y+1}$

ii) $(3xy + 8y^5)dx + (2x^2 + 24xy^4)dy = 0$

iii) $\frac{dy}{dx} = \frac{y}{2y \log y + y - x}$

OR

Q2) a) Form the differential equation whose general solution is
 $y = (C_1 + C_2 t)e^t$ [6]

b) Solve any two [10]

i) $\frac{dy}{dx} = 1 - x \tan(x - y)$

ii) $(xy - 2y^2)dx - (x^2 - 3xy)dy = 0$

iii) $(x + \tan y)dy = \sin 2y dx$

P.T.O

Q3) Solve any three:

[18]

- a) Radium decomposes at the rate proportional to the amount present. If 5% of the original amount disappears in 50 years, how much will remain after 100 years.
- b) A constant e.m.f E volts is applied to a circuit containing a constant resistance R ohms in series and a constant inductance L henries. If the initial current is zero, show that the current builds up to half its theoretical maximum in $\left(\frac{L \log 2}{R}\right)$ seconds
- c) A body at temperature 100°C is placed in a room whose temperature is 20°C and cools to 60°C in 5 minutes. Find its temperature after 10 minutes.
- d) Find the orthogonal trajectories of $r = a(1 - \cos \theta)$

OR

Q4) Solve any three:

[18]

- a) The inner and outer surfaces of a spherical shell are maintained at T_0 and T_1 temperatures respectively. If the inner and outer radii are r_0 and r_1 respectively and thermal conductivity of the shell is k , Find the amount of heat lost from the shell per unit time. Find also the temperature distribution through the shell.
- b) A body starts moving from rest and is opposed by a force per unit mass of value cx and resistance per unit mass of value bv^2 , where x is displacement and v is velocity of the body at that instant. show that the velocity of the particle is given by $v^2 = \frac{c}{2b^2} (1 - e^{-2bx}) - \frac{cx}{b}$
- c) If the population of a country doubles in 50 years in how many years will it become three times under the assumption that the rate of increase is proportional to the number of inhabitants ?

- d) A body originally at 80° C cools down to 60° C in 20 minutes, the temperature of air being 40° C. What will be temperature of the body after 80 minutes from the original?

- Q5)** a) Find the Fourier series to represent $f(x) = x^2$ in the interval $-l < x < l$. [9]

Hence deduce:

$$\frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$$

- b) Find the reduction formula for, [7]

$$I_n = \int_0^{\infty} e^{-x} \sin^n x dx$$

Hence show : $I_4 = \frac{24}{85}$

OR

- Q6)** a) Express 'y' as a Fourier series upto first harmonic where y is given as: [8]

x	0	$\frac{\pi}{3}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{5\pi}{3}$
y	1.0	1.4	1.9	1.7	1.5	1.2

- b) Evaluate : $\int_0^{\infty} \frac{x^5}{5^x} dx$ [4]

- c) Prove that : [4]

$$B(m, n) = B(m, n + 1) + B(m + 1, n)$$

SECTION - II

Q7) a) Trace the following curves (any two) **[8]**

i) $x(x^2 + y^2) = a(x^2 - y^2) a > 0$

ii) $r = a \sin 2\theta$

iii) $x = a(t + \sin t), y = a(1 - \cos t)$

b) Find the length of one loop of lemniscate **[5]**

$$r^2 = a^2 \cos 2\theta$$

c) if $\phi(a) = \int_a^{a^2} \frac{\sin ax}{x} dx$ find $\frac{d\phi}{da}$ **[4]**

OR

Q8) a) Trace the following curves (any two) **[8]**

i) $a^2 y^2 = x^2 (a^2 - x^2)$

ii) $r = \frac{a}{2}(1 - \sin \theta)$

iii) $x^{2/3} + y^{2/3} = a^{2/3}$

b) Show that $\frac{d}{dt} \operatorname{erf} \sqrt{t} = \frac{e^{-t}}{\sqrt{\pi t}}$ **[4]**

c) In Astroid $x^{2/3} + y^{2/3} = a^{2/3}$, arc length S is measured from cusp, which lies on y - axis, Show that $S^3 \propto x^2$ **[5]**

Q9) a) Find the equation of circle which is the section of sphere $x^2 + y^2 + z^2 + 6y - 6z - 21 = 0$ and has center on $M(2, -1, 2)$ [6]

b) Find the equation of right circular cone with vertex $(1, 2, -3)$ and semivertical angle $\cos^{-1} \frac{1}{\sqrt{3}}$, whose axis is $\frac{x-1}{1} = \frac{y-2}{2} = \frac{z+1}{-1}$ [5]

c) Find the equation of right circular cylinder with radius 5 and axis is the line $\frac{x-2}{2} = \frac{y-3}{1} = \frac{z+1}{1}$ [6]

OR

Q10) a) A sphere with constant radius r passes through origin and meets co-ordinate axes in A, B, C respectively, Show that the locus of centroid of the triangle ABC is a Sphere $\mathcal{S} (x^2 + y^2 + z^2) = 4r^2$ [6]

b) Find the equation of right circular cone passing through $(1, 1, 2)$ with vertex as origin and axis is $6x = -3y = 4z$. [5]

c) Find the equation of right circular cylinder with axis [6]

$$\frac{x-1}{2} = \frac{y-2}{1} = \frac{z-3}{2} \text{ and radius} = 2$$

Q11) Solve any two [16]

a) Evaluate $\int_0^a \int_0^{\sqrt{a^2-x^2}} e^{-x^2-y^2} dx dy$

b) Prove that the mean distance of the points within a circular area of radius a , from a fixed point on the circumference is $\frac{32a}{9\pi}$

c) Find the volume of the region bounded by $z = x^2 + y^2$ and $z = 2x$

OR

Q12) Solve any two

[16]

a) Evaluate $\iiint_V \frac{dx dy dz}{(1+x+y+z)^3}$

Over the volume of tetrahedron bounded by

$$x=0, y=0, z=0 \text{ and } x+y+z=1$$

b) Find the centre of Gravity of the area in first quadrant, bounded by

$$\left(\frac{x}{a}\right)^{\frac{2}{3}} + \left(\frac{y}{b}\right)^{\frac{2}{3}} = 1$$

Where density $\rho = \lambda xy, \lambda = \text{constant}$

c) Find the area between the parabola $\frac{y+8}{x} = x-2$ and X – axis.

