

Total No. of Questions—12]

[Total No. of Printed Pages—8

<b>Seat No.</b>	
---------------------	--

**[4756]-21**

**F.E. (Second Semester) EXAMINATION, 2015**

**ENGINEERING MATHEMATICS-II**

**(2008 PATTERN)**

**Time : Three Hours**

**Maximum Marks : 100**

**N.B. :— (i) Section I : Solve Q. No. 1 or Q. No. 2, Q. No. 3 or Q. No. 4, Q. No. 5 or Q. No. 6.**

**Section II : Solve Q. No. 7 or Q. No. 8, Q. No. 9 or Q. No. 10, Q. No. 11 or Q. No. 12.**

- (ii) Neat diagrams must be drawn wherever necessary.
- (iii) Figures to the right indicate full marks.
- (iv) Use of non-programmable electronic pocket calculator is allowed.
- (v) Assume suitable data, if necessary.

**SECTION I**

1. (a) Form the differential equation whose general solution is :

$$y = e^x [c_1 \cos x + c_2 \sin x],$$

where  $c_1$  and  $c_2$  are arbitrary constants.

[6]

P.T.O.

(b) Solve any *two* : [10]

(i)  $\frac{dy}{dx} = (4x + y)^2$

(ii)  $(x^2 - 3xy + 2y^2) dx + (3x^2 - 2xy) dy = 0$

(iii)  $\frac{dy}{dx} + \frac{4x}{1+x^2} y = \frac{1}{(x^2+1)^3}$

Or

2. (a) Form the differential equation by eliminating arbitrary constants  $c_1$  and  $c_2$  from the general solution given by : [6]

$$y = c_1 \cos \log x + c_2 \sin \log x$$

(b) Solve any *two* : [10]

(i)  $\frac{dy}{dx} = \frac{6x - 4y + 3}{3x - 2y + 1}$

(ii)  $\frac{dy}{dx} - \frac{\tan y}{1+x} = (1+x) e^x \sec y$

(iii)  $(2x + e^x \log y) y dx + e^x dy = 0.$

3. Solve any *three* : [18]

(a) Find the orthogonal trajectories of the family :

$$xy = c^2$$

- (b) An e.m.f.  $200 e^{-5t}$  is applied to a series circuit consisting of  $20 \Omega$  resistor and  $0.01 \text{ F}$  capacitor. Find the charge and current at any time, assuming that there is no initial voltage on capacitor.
- (c) A body of mass  $m$  falling from a rest is subject to the force of gravity and an air resistance proportional to the square of velocity ( $kv^2$ ). If it falls through a distance  $x$  and possesses a velocity  $v$  at that instant. Prove that :

$$\frac{2 kx}{m} = \log \left( \frac{a^2}{a^2 - v^2} \right)$$

where  $mg = ka^2$ .

- (d) If 30% of radioactive substance disappeared in 10 days, how long will it take for 90% of it to disappeared ?

*Or*

4. Solve any *three* : [18]

- (a) A body of temperature  $100^\circ\text{C}$  is placed in a room whose temperature is  $20^\circ\text{C}$  and cools to  $60^\circ\text{C}$  in 5 minutes. What is time required to reach temperature of body at  $40^\circ\text{C}$ .

- (b) A voltage  $10 e^{-2t}$  is applied at  $t = 0$  to a circuit containing an inductance  $L$  and resistance  $R$  connected in a series. Find current  $I$  at any time  $t$  as a function of time  $t$ , given that when  $t = 0$ ,  $I = 0$ .
- (c) A long hollow pipe has an inner diameter of 10 cm and outer diameter of 20 cm, the inner surface is kept at  $200^\circ\text{C}$  and outer surface is at  $50^\circ\text{C}$ . The thermal conductivity  $k = 0.12$ . How much heat is lost per minute from the portion of the pipe 20 m long.
- (d) A metal ball is heated to a temperature of  $100^\circ\text{C}$  at time  $t = 0$ , it is placed in a water which is maintained at  $40^\circ\text{C}$ . If the temperature of the ball reduces to  $60^\circ\text{C}$  in 4 minutes, find the time at which the temperature of ball is  $50^\circ\text{C}$

5. (a) Find the Fourier series for :

$$f(x) = x^2, \quad -\pi < x < \pi$$

and hence deduce that : [9]

$$\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots = \frac{\pi^2}{12}$$

(b) If :

$$U_n = \int_0^{\pi/4} \tan^n \theta \, d\theta$$

show that :

$$n (U_{n+1} + U_{n-1}) = 1$$

hence find  $U_4$ . [7]

*Or*

6. (a) The following table gives variation of periodic current over a period : [4]

<b>t (sec)</b>	:	0	T/6	T/3	T/2	2T/3	$\frac{5T}{6}$	T
<b>A (amp)</b>	:	1.98	1.30	1.05	1.30	-0.88	-0.25	1.98

Show that there is a direct current part of 0.75 amp in variable current and obtain the amplitude of first harmonic. [8]

(b) Evaluate : [4]

$$\int_0^{\infty} e^{-2x^2} x^9 \, dx$$

(c) Prove that : [4]

$$\beta(m, n) = \int_0^1 \frac{x^{m-1} + x^{n-1}}{(1+x)^{m+n}} \, dx.$$

## SECTION II

7. (a) Trace the following curves (any two) : [8]

(i)  $xy^2 = a^2 (2a - x)$

(ii)  $r = a(1 - \sin \theta)$

(iii)  $x^{2/3} + y^{2/3} = a^{2/3}$

(b) Show that :

$$\phi(a) = \int_{\pi/6a}^{\pi/2a} \frac{\sin ax}{x} dx$$

is independent of  $a$ . [4]

(c) Find the perimeter of the cardioid : [5]

$$r = a (1 - \cos \theta)$$

*Or*

8. (a) Trace the following curves (any two) : [8]

(i)  $y^2 (3a - x) = x^3$

(ii)  $r = a \sin 2\theta$

(iii)  $x = t^2, y = t - \frac{t^3}{3}$

(b) Show that : [4]

$$\int_0^{\infty} e^{-x^2 - 2ax} dx = \frac{\sqrt{\pi}}{2} e^{a^2} [1 - \operatorname{erf}(a)]$$

(c) Find the whole length of the loop of the curve : [5]

$$3y^2 = x(x - 1)^2$$

9. (a) Prove that the sphere :

$$x^2 + y^2 + z^2 + 2x - 4y - 2z - 3 = 0$$

touches the plane :

$$2x - 2y - z + 16 = 0$$

and find the point of contact.

- (b) Find the equation of the right circular cone whose vertex is at the origin, axis is the line :

$$\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$$

and semi-vertical angle is  $30^\circ$ . [5]

- (c) Find the equation of the right circular cylinder with radius 2 and axis is the line : [6]

$$\frac{x - 1}{2} = \frac{y - 2}{1} = \frac{z - 3}{2}$$

*Or*

10. (a) Find the equation of the sphere which has its centre at (2, 3, -1) and which touches the line : [6]

$$\frac{x + 1}{-5} = \frac{y - 8}{3} = \frac{z - 4}{4}$$

- (b) Find the equation of the right circular cone with vertex (1, 2, 3) axis has direction ratios 2, -1, 4 and semi-vertical angle is  $60^\circ$ . [5]

(c) Find the equation of the right circular cylinder with axis :

$$\frac{x-1}{2} = \frac{y-3}{2} = \frac{z-5}{-1}$$

and radius = 3. [6]

11. Solve any two :

(a) Evaluate : [8]

$$\int_0^1 \int_0^{\sqrt{1+x^2}} \frac{dy dx}{(1+x^2+y^2)}$$

(b) Find the area of the upper half of the cardioid : [8]

$$r = a(1 + \cos \theta)$$

(c) Find the C.G of an arc of the catenary : [8]

$$y = a \cosh\left(\frac{x}{a}\right) \text{ from } x = -a \text{ to } x = a.$$

Or

12. Solve any two :

(a) Evaluate : [8]

$$\int_0^1 \int_0^{\sqrt{1-x^2}} \frac{y dy dx}{(1+y^2)\sqrt{1-x^2-y^2}}$$

(b) Evaluate : [8]

$$\int_0^{\log 2} \int_0^x \int_0^{x+y} e^{x+y+z} dx dy dz$$

(c) Find the moment of inertia of a sphere about a diameter. [8]