

Total No. of Questions—8]

[Total No. of Printed Pages—7

| | |
|-------------|--|
| Seat No. | |
|-------------|--|

[5352]-505

S.E. (Civil) (I Sem.) EXAMINATION, 2018

ENGINEERING MATHEMATICS—III

(2015 PATTERN)

Time : Two Hours

Maximum Marks : 50

N.B. :— (i) Answer Q. No. 1 or Q. No. 2, Q. No. 3 or Q. No. 4,
Q. No. 5 or Q. No. 6, Q. No. 7 or Q. No. 8.

(ii) Draw neat diagrams wherever needed.

(iii) Figures to the right indicate full marks.

(iv) Assume suitable data, if necessary.

(v) Use of non-programmable pocket calculator is allowed.

1. (a) Solve any *two* of the following : [8]

(i) $(D^3 - D^2 + 4D - 4) y = e^x$

(ii) $(D^2 - 2D + 2) y = e^x \tan x$

(by method of variation of parameters)

(iii) $x^3 \frac{d^3 y}{dx^3} + x^2 \frac{d^2 y}{dx^2} - 2y = \frac{1}{x^3}$.

(b) Apply Gauss-Jordan method to solve the equations : [4]

$$x + y + z = 9$$

$$2x - 3y + 4z = 13$$

$$3x + 4y + 5z = 40.$$

P.T.O.

Or

2. (a) A light horizontal strut AB of length l is freely pinned at A & B and is under the action of equal and opposite compressive forces P at each of its ends and carries a load W at its centre. Show that the deflection at its centre is : [4]

$$\frac{W}{2P} \left[\frac{1}{n} \tan \frac{nl}{2} - \frac{l}{2} \right] \text{ where } n^2 = \frac{P}{EI}.$$

- (b) Using fourth order Runge-Kutta method solve the equation : [4]

$$\frac{dy}{dx} = \sqrt{x + y}$$

subject to the conditions $x = 0, y = 1$ to find y at $x = 0.1$ taking $h = 0.1$.

- (c) Solve the following system by Cholesky's method : [4]

$$9x_1 + 6x_2 + 12x_3 = 17.4$$

$$6x_1 + 13x_2 + 11x_3 = 23.6$$

$$12x_1 + 11x_2 + 26x_3 = 30.8.$$

3. (a) The equation of two lines of regression obtained in a correlation analysis are the following : [4]

$$2x + 3y - 8 = 0 \text{ and}$$

$$x + 2y - 5 = 0.$$

Obtain the value of the correlation co-efficient and the variance of y given that variance of x is 12.

(b) An aptitude test for selecting officers in a bank conducted on 1000 candidates. The average score is 42 and standard deviation of score is 24. Assuming normal distribution for the score find :

- (i) The number of candidates whose scores exceed 60.
(ii) The number of candidates whose score lie between 30 and 60.

[Given Area = 0.2734 for $z = 0.75$, Area = 0.1915 for $z = 0.5$.]

(c) Find the directional derivative of $\phi = 4e^{2x - y + z}$ at the point (1, 1, -1) in the direction towards (-3, 5, 6). [4]

Or

4. (a) In a certain distribution the first four moments about 4, are 1.5, 17, 30 and 108. Find the central moments and hence β_1 and β_2 . [4]

(b) Prove the following (any one) : [4]

(i)
$$\nabla \left(\frac{\bar{a} \cdot \bar{r}}{r^n} \right) = \frac{\bar{a}}{r^n} - \frac{n(\bar{a} \cdot \bar{r})}{r^{n+2}} \bar{r}$$

(ii)
$$\bar{a} \cdot \nabla \left[\bar{b} \cdot \nabla \left(\frac{1}{r} \right) \right] = \frac{3(\bar{a} \cdot \bar{r})(\bar{b} \cdot \bar{r})}{r^5} - \frac{\bar{a} \cdot \bar{b}}{r^3}$$

(c) Show that the vector field : [4]

$$\bar{F} = (y^2 \cos x + z^2) \hat{i} + (2y \sin x) \hat{j} + 2xz \hat{k}$$

is conservative and find scalar field such that $\bar{F} = \nabla \phi$.

5. Attempt any two :

(a) Using Green's theorem evaluate : [6]

$$\oint \bar{F} \cdot d\bar{r}$$

for the field

$$\bar{F} = 2x^2y\bar{i} + x^3\bar{j}$$

over the first quadrant of the circle $x^2 + y^2 = a^2$.

(b) Using Divergence theorem evaluate : [6]

$$\iiint_S (y^2z^2\bar{i} + z^2x^2\bar{j} + x^2y^2\bar{k}) \cdot d\bar{S},$$

where S is the upper half of sphere $x^2 + y^2 + z^2 = 4$ above the plane $z = 0$.

(c) Evaluate : [7]

$$\iiint_S \nabla \times \bar{F} \cdot \hat{n} dS,$$

where

$$\bar{F} = (x - y)\bar{i} + (x^2 + yz)\bar{j} - 3xy^2\bar{k}$$

and S is the surface of the cone $z = 4 - \sqrt{x^2 + y^2}$ above the XOY-plane.

Or

6. Attempt any two :

(a) Find the work done in moving a particle along $x = 3 \cos \theta$,

$y = 3 \sin \theta$, $z = 5\theta$, from $\theta = \frac{\pi}{4}$ to $\theta = \frac{\pi}{2}$ under a field

of force given by : [6]

$$\bar{F} = -9\sin^2\theta\cos\theta\bar{i} + 3(2\sin\theta - 3\sin^3\theta)\bar{j} + 5\sin 2\theta\bar{k}.$$

(b) Evaluate : [6]

$$\iint_S (x \bar{i} + y \bar{j} + z \bar{k}) \cdot d\bar{S}$$

where S is the curved surface of the cylinder $x^2 + y^2 = 4$ bounded by the planes $z = 0$ and $z = 2$.

(c) Evaluate : [7]

$$\iint_S (\nabla \times \bar{F}) \cdot d\bar{S}$$

where

$$\bar{F} = x^3 \bar{i} - xyz \bar{j} + y^3 \bar{k}$$

and S is the surface $x^2 + 9y^2 + 4z^2 - 2x = 36$ above the plane $x = 0$.

7. Solve any *two* of the following :

(a) If $\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$ represents the vibrations of a string of length l fixed at both ends find the solution with the following conditions : [7]

(i) $y(0, t) = 0$

(ii) $y(l, t) = 0$ for all t ,

(iii) $\left(\frac{\partial y}{\partial t}\right)_{t=0} = 0$ for all x and

(iv) $y(x, 0) = \frac{3a}{2l}x, \quad 0 \leq x \leq \frac{2l}{3}$
 $= \frac{3a}{l}(l - x), \quad \frac{2l}{3} \leq x \leq l.$

- (b) A rod of length l with insulated sides is initially at a uniform temperature x . Both the ends of the rod are kept at zero temperature. Find the temperature at any point and at any time t , use $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$. [6]
- (c) A rectangular plate is bounded by $x = 0$, $x = a$, $y = 0$, $y = b$. Its surfaces are insulated and temperature along three edges $x = 0$, $x = a$, $y = 0$ is maintained at 0°C , while the fourth edge $y = b$ is maintained at constant temperature u_0 until steady state is reached. Find steady state temperature $u(x, y)$. [6]

Or

8. Solve any *two* of the following :

- (a) If $\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$ represents the vibrations of string of length l fixed at both ends, find the solution with boundary conditions : [7]
- (i) $y(0, t) = 0, \quad \forall t$
- (ii) $y(l, t) = 0, \quad \forall t$ and initial conditions
- (iii) $\left(\frac{\partial y}{\partial t}\right)_{t=0} = 0, \quad \forall x$
- (iv) $y(x, 0) = k(lx, x^2), \quad 0 \leq x \leq l$

(b) The temperature at any point of a insulated metal rod of one meter length is governed by the differential equation $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$. Find $u(x, t)$ subject to the following conditions : [6]

(i) $u(0, t) = 0^\circ\text{C}$

(ii) $u(1, t) = 0^\circ\text{C}$

(iii) $u(x, 0) = 50^\circ\text{C}$,

and hence find the temperature in the middle of the rod at subsequent time.

(c) Solve $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$, which satisfies the conditions : [6]

$u(0, y) = u(l, y) = u(x, 0) = 0$ and

$u(x, a) = \sin \frac{m\pi}{l} x, \quad 0 \leq x \leq l.$